Author Index

Abe, S., see Yamada, K. 367
Achari, R.
–, Drissel, D., Thomas, D., Shin, K. and Look, Z.
Analysis of esmolol in human blood by high-performance liquid chromatography and its application to pharmacokinetic studies 430
Adlard, M.W., see Shah, A.J. 325
Akane, A., see Matsubara, K. 49
Alakhov, Y.B., see Prisyazhnoy, V.S. 243
Alfredsson, G.
–, Wiesel, F.-A. and Lindberg, M.
Glutamate and glutamine in cerebrospinal fluid and serum from healthy volunteers - analytical aspects 378
Anderson, G.M.
–, Durkin, T.A., Morton, J.B. and Cohen, D.J.
Liquid chromatographic determination of urinary catecholamines after one-step alumina extraction 373
Argoudelis, C.J.
Simple isocratic high-performance liquid chromatographic method for the separation of the six vitamers of vitamin B_6 315
Arnoux, Ph., see Payet, B. 337
Baars, A.M., see Vree, T.B. 440
Beightol, L.A., see Karnes, H.T. 398
Benfenati, E., see Latini, R. 211
Bochner, F., see Chen, Z.R. 170
Booij, L.H.D.J., see Vree, T.B. 440
Bos, O.J.M.
–, Fischer, M.J.E., Wilting, J. and Janssen, L.H.M.
Procedure for the isolation of a large peptic fragment of human serum albumin 13
Bourne, D.W.A., see Hakim, L. 111
Brain, K.R., see Pourgholami, M.H. 163
Brinkman, U.A.Th., see Irth, H. 95
Bromet-Petit, M., see Duchene, P. 205
Bucheli, F., see Wyss, R. 303
Cano, J.P., see Payet, B. 337
Carlucci, G., see Palumbo, G. 201
Catalin, J., see Payet, B. 337
Caude, M., see Ni, P. 255
Chen, Z., see Latini, R. 211
Chen, Z.R.
–, Siebert, D.M., Somogyi, A.A. and Bochner, F.
Determination of pholcodine in biological fluids by high-performance liquid chromatography with fluorescence detection 170
Cohen, D.J., see Anderson, G.M. 373
Corre, P. Le, see Le Corre, P. 424
Corti, F.
–, Luzzani, F. and Ventura, P.
Sulphate quantification in rabbit tracheobronchial mucus by reversed-phase ion-pair chromatography 147
Cox, J.W.
– and Pullen, R.H.
Irregular retention properties of 21-aminosteroid antioxidants in octylsilane bonded-phase high-performance liquid chromatography 285
– and Pullen, R.H.
High-performance liquid chromatographic determination of a 21-aminosteroid antioxidant in plasma 293
Czejka, M.J.
– and Georgopoulos, A.
High-performance liquid chromatographic determination of adriblastin in human plasma, urine, saliva and liver punctuate by column switching for drug monitoring studies 182
Dale, G.L.
Use of the dye Janus Green to facilitate quantitative analysis of phosphoinositides 445
De Jong, G.J., see Irth, H. 95
Dily, J. le, see Duchene, P. 205
Drissel, D., see Achari, R. 430
Duchene, P.
–, Le Dily, J., Bromet-Petit, M., Mosser, J. and Feniou, C.
High-performance liquid chromatographic assay of the metabolites of thymoxamine 205
Durkin, T.A., see Anderson, G.M. 373
Eckerlin, R.H., see Hsu, S.-H. 219
Edholm, L.-E.
–, Lindberg, C., Paulson, J. and Walhagen, A.
Determination of drug enantiomers in biological samples by coupled column liquid chromatography and liquid chromatography–mass spectrometry 61
Ellis, P.P., see Riegel, M.A. 177
Emilsson, H.
High-performance liquid chromatographic determination of cadralazine in human plasma and urine 119

Farthing, D., see Karnes, H.T. 398
Feniou, C., see Duchene, P. 205
Fischer, M.J.E., see Bos, O.J.M. 13
Förster, H.-G.
–, Siems, W., Rathmann, J. and Müller, M.
New method of purification of perchloric acid extracts for quantitative gas chromatography of some Krebs cycle acids 357
Frei, R.W., see Irth, H. 95
Fukasawa, K., see Harada, M. 129
Fukasawa, K.M., see Harada, M. 129
Fukui, Y., see Matsubara, K. 49
Fusek, M., see Prisyazhnoy, V.S. 243
Gaida, A.V.
–, Monastyrskii, V.A., Magerovskii, Yu.V., Staroverov, S.M. and Lisichkin, G.V.
Affinity chromatography of human thrombin on modified silica 385
Gavinelli, M., see Latini, R. 211
Georgopoulos, A., see Czejka, M.J. 182
Gibassier, D., see Le Corre, P. 424
Greenblatt, D.J., see Locniskar, A. 215
Gupta, M.B.
–, Hubbard, J.W. and Midha, K.K.
Separation of enantiomers of derivatised or underivatised propranolol by means of high-performance liqud chromatography 189
Guyon, F., see Ni, P. 255
Hakim, L.
–, Bourne, D.W.A. and Triggs, E.J.
High-performance liquid chromatographic assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 111
Harada, M.
–, Hiraoka, B.Y., Mogi, M., Fukasawa, K. and Fukasawa, K.M.
High-performance liquid chromatographic separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 129
Hart, B.J. 't, see 't Hart, B.J. 403
Hashida, N., see Seki, T. 410
Hashimoto, Y., see Yamashita, K. 39
Henion, J.D., see Hsu, S.-H. 219
Heykants, J., see Woestenborghs, R. 195
Hiraoka, B.Y., see Harada, M. 129
Holthuis, J.J.M., see Vendrig, D.E.M.M. 83
Honkasalo, T.
– and Nissinen, E.
Determination of phenol sulphotransferase activity by high-performance liquid chromatography 136
Howell, S.B., see Velury, S. 141
Hsu, S.-H.
–, Eckerlin, R.H. and Henion, J.D.
Identification and quantitation of trenbolone in bovine tissue by gas chromatography–mass spectrometry 219
Hubbard, J.W., see Gupta, M.B. 189
Irth, H.
–, De Jong, G.J., Brinkman, U.A.Th. and Frei, R.W.
Determination of disulfiram and two of its metabolites in urine by reversed-phase liquid chromatography and spectrophotometric detection after post-column complexation 95
Ishibashi, M., see Yamashita, K. 1
Iwai, K., see Shibata, K. 23
Jaeger, H., see Nieder, M. 73
Janssen, L.H.M., see Bos, O.J.M. 13
Jong, G.J. de, see Irth, H. 95
Kagawa, M., see Matsubara, K. 49
Kanazawa, T., see Seki, T. 410
Kaneda, N., see Sasaoka, T. 392
Karle, J.M.
– and Olmeda, R.
Rapid and sensitive quantitative analysis of the new antimalarial N^4-[2,6-dimethoxy-4-methyl-5-[(3-trifluoromethyl)phenoxy]-8-quinolinyl]-1,4-pentanediamine in plasma by liquid chromatography and electrochemical detection 347
Karnes, H.T.
–, Beightol, L.A., Serafin, R.J. and Farthing, D.
Improved method for the determination of diazepam and N-desmethyldiazepam in plasma using capillary gas chromatography and nitrogen–phosphorus detection 398
Katayama, K., see Yamada, K. 367
Katori, M., see Yamashita, K. 1
Kawada, T., see Shibata, K. 23
Kawanabe, K., see Tanabe, S. 29
Kitahara, M., see Tanabe, S. 29
Kitazawa, Y., see Takeyama, N. 361
Knijff, J.H., see Odink, J. 273
Knudsen, J.J.
– and McGown, E.L.
Gas chromatographic analysis of urinary dimercaptosuccinic acid 231
Koga, R., see Yamashita, K. 39
Korthals, H., see Odink, J. 273
Latini, R.
–, Sica, A., Marchi, S., Chen, Z., Gavinelli, M. and Benfenati, E.
High-performance liquid chromatographic

separation and mass spectrometric identification of propafenone, 5-hydroxypropafenone and N-depropylpropafenone 211
Le Corre, P.
–, Gibassier, D., Sado, P. and Le Verge, R.
Simultaneous assay of disopyramide and monodesisopropyldisopyramide enantiomers in biological samples by liquid chromatography 424
Le Dily, J., see Duchene, P. 205
Le Verge, R., see Le Corre, P. 424
Lindberg, C., see Edholm, L.-E. 61
Lindberg, M., see Alfredsson, G. 378
Lisichkin, G.V., see Gaida, A.V. 385
Locniskar, A.
– and Greenblatt, D.J.
High-performance liquid chromatographic assay of adinazolam and its desmethyl metabolite 215
Look, Z., see Achari, R. 430
Lorreyne, W., see Woestenborghs, R. 195
Luzzani, F., see Corti, F. 147
McGown, E.L., see Knudsen, J.J. 231
Maessen, P.A.
–, Pinedo, H.M., Mross, K.B. and Van der Vijgh, W.J.F.
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
Magerovskii, Yu.V., see Gaida, A.V. 385
Marchi, S., see Latini, R. 211
Maseda, C., see Matsubara, K. 49
Matsubara, K.
–, Akane, A., Takahashi, S., Shiono, H., Fukui, Y., Kagawa, M. and Maseda, C.
Gas chromatographic determination for forensic purposes of petroleum fuel inhaled just before fatal burning 49
Mellström, B.
High-performance liquid chromatographic quantitation of 4-hydroxydebrisoquine in microsomal incubates by use of silica columns, aqueous mobile phase and automated column switching 435
Midha, K.K., see Gupta, M.B. 189
Miyazaki, H., see Yamashita, K. 1
Mizuguchi, S., see Yamashita, K. 39
Mogi, M., see Harada, M. 129
Molinaro, N.C., see Smith, H.T. 416
Monastyrskii, V.A., see Gaida, A.V. 385
Morton, J.B., see Anderson, G.M. 373
Mosser, J., see Duchene, P. 205
Mross, K.B., see Maessen, P.A. 103
Müller, M., see Förster, H.-G. 357
Nagatsu, T., see Sasaoka, T. 392
Nakamura, T., see Yamashita, K. 39
Nawata, M., see Tanabe, S. 29
Ni, P.
–, Guyon, F., Caude, M. and Rosset, R.
Automatic determination of urinary 4-hydroxy-3-methoxymandelic (vanillylmandelic) acid by liquid chromatography with electrochemical detection 255
Nicholls, P.J., see Pourgholami, M.H. 163
Nieder, M.
– and Jaeger, H.
Sensitive quantification of pseudoephedrine in human plasma and urine by high-performance liquid chromatography 73
Nissinen, E., see Honkasalo, T. 136
Odink, J.
–, Korthals, H. and Knijff, J.H.
Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by liquid chromatography with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 273
Olmeda, R., see Karle, J.M. 347
Oss, G.E.C.J.M. van, see Vree, T.B. 440
Palumbo, G.
–, Pantaleoni, G.C. and Carlucci, G.
Determination of furprofen in human plasma by solid-phase extraction and reversed-phase high-performance liquid chromatography 201
Pantaleoni, G.C., see Palumbo, G. 201
Paterson, S.
Measurement of dipipanone using capillary gas chromatography 152
Paulson, J., see Edholm, L.-E. 61
Payet, B.
–, Arnoux, Ph., Catalin, J. and Cano, J.P.
Direct determination of mitoxantrone and its mono- and dicarboxylic metabolites in plasma and urine by high-performance liquid chromatography 337
Peaston, R.T.
Routine determination of urinary free catecholamines by high-performance liquid chromatography with electrochemical detection 263
Pinedo, H.M., see Maessen, P.A. 103
Pourgholami, M.H.
–, Brian, K.R. and Nicholls, P.J.
High-performance liquid chromatographic determination of imide inhibitors of aro-

matase in biological samples 163
Prisyazhnoy, V.S.
–, Fusek, M. and Alakhov, Y.B.
Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their $F_{ab'}$ fragments for isolation of proteins 243
Pullen, R.H., see Cox, J.W. 285, 293
Rathmann, J., see Förster, H.-G. 357
Riegel, M.A.
– and Ellis, P.P.
High-performance liquid chromatographic assay for piperacillin in aqueous humor of the eye 177
Rompaey, F. Van, see Woestenborghs, R. 195
Rosset, R., see Ni, P. 255
Sado, P., see Le Corre, P. 424
Sasaoka, T.
–, Kaneda, N. and Nagatsu, T.
Highly sensitive assay for xanthine oxidase activity by high-performance liquid chromatography with fluorescence detection 392
Sato, T., see Yamada, K. 367
Seki, T.
–, Hashida, N. and Kanazawa, T.
Determination of tetraprenylacetone in human plasma by high-performance liquid chromatography with fluorescence derivatization using dansylhydrazine 410
Serafin, R.J., see Karnes, H.T. 398
Shah, A.J.
– and Adlard, M.W.
Determination of β-lactams and their biosynthetic intermediates in fermentation media by pre-column derivatization followed by fluorescence detection 325
Shibata, K.
–, Kawada, T. and Iwai, K.
Simultaneous micro-determination of nicotinamide and its major metabolites, N^1-methyl-2-pyridone-5-carboxamide and N^1-methyl-4-pyridone-3-carboxamide, by high-performance liquid chromatography 23
Shin, K., see Achari, R. 430
Shiono, H., see Matsubara, K. 49
Sica, A., see Latini, R. 211
Siebert, D.M., see Chen, Z.R. 170
Siems, W., see Förster, H.-G. 357
Smith, H.T.
– and Molinaro, N.C.
High-performance liquid chromatographic method for the determination of methysergide and methylergonovine in human plasma 416
Somogyi, A.A., see Chen, Z.R. 170
Staroverov, S.M., see Gaida, A.V. 385
Stiffin, R.M., see Wainer, I.W. 158
Takagi, D., see Takeyama, N. 361
Takahashi, S., see Matsubara, K. 49
Takeyama, N.
–, Takagi, D., Kitazawa, Y. and Tanaka, T.
Determination of α-keto acids in rat hindquarter muscle by high-performance liquid chromatography with fluorescence detection 361
Tanabe, S.
–, Kitahara, M., Nawata, M. and Kawanabe, K.
Determination of oxidizable inorganic anions by high-performance liquid chromatography with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 29
Tanaka, T., see Takeyama, N. 361
Teeuwsen, J., see Vendrig, D.E.M.M. 83
't Hart, B.J.
– and Wilting, J.
Sensitive gas chromatographic method for determining nitrazepam in serum and saliva 403
Thomas, D., see Achari, R. 430
Triggs, E.J., see Hakim, L. 111
Umezawa, H., see Yamashita, K. 39
Van der Vijgh, W.J.F., see Maessen, P.A. 103
Van Oss, G.E.C.J.M., see Vree, T.B. 440
Van Rompaey, F., see Woestenborghs, R. 195
Velury, S.
– and Howell, S.B.
Measurement of plasma thiols after derivatization with monobromobimane 141
Vendrig, D.E.M.M.
–, Teeuwsen, J. and Holthuis, J.J.M.
Analysis of vinca alkaloids in plasma and urine using high-performance liquid chromatography with electrochemical detection 83
Ventura, P., see Corti, F. 147
Verge, R. Le, see Le Corre, P. 424
Vijgh, W.J.F. van der, see Maessen, P.A. 103
Vree, T.B.
–, Baars, A.M., Van Oss, G.E.C.J.M. and Booij, L.H.D.J.
High-performance liquid chromatography and preliminary pharmacokinetics of articaine and its 2-carboxy metabolite in human serum and urine 440

Wainer, I.W.
– and Stiffin, R.M.
Direct resolution of the stereoisomers of leucovorin and 5-methyltetrahydrofolate using a bovine serum albumin high-performance liquid chromatographic chiral stationary phase coupled to an achiral phenyl column 158
Walhagen, A., see Edholm, L.-E. 61
Watanabe, K., see Yamashita, K. 1, 39
Wiesel, F.-A., see Alfredsson, G. 378
Wilting, J., see Bos, O.J.M. 13
–, see 't Hart, B.J. 403
Woestenborghs, R.
–, Lorreyne, W., Van Rompaey, F. and Heykants, J.
Determination of cisapride in plasma and animal tissues by high-performance liquid chromatography 195
Wyss, R.
– and Bucheli, F.
Quantitative analysis of retinoids in biological fluids by high-performance liquid chromatography using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo -metabolites in plasma 303
Yamada, K.
–, Abe, S., Katayama, K. and Sato, T.
Sensitive high-performance liquid chromatographic method for the determination of phosphatidic acid 367
Yamashita, K.
–, Watanabe, K., Ishibashi, M., Katori, M. and Miyazaki, H.
Dimethylisopropylsilyl ether derivatives in gas chromatography–mass spectrometry of some prostanoids. Application to the microdetermination of 15-ketoprostaglandin $F_{2\alpha}$ formed by enzymatic conversion of $PGF_{2\alpha}$ in vitro 1
–, Watanabe, K., Koga, R., Mizuguchi, S., Hashimoto, Y., Nakamura, T. and Umezawa, H.
Determination of deoxyspergualin by gas chromatography–selected-ion monitoring 39

Subject Index

Adinazolam
HPLC assay for adinazolam and its desmethyl metabolite 215

Adriamycin
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103

Adriblastin
HPLC determination of adriblastin in human plasma, urine saliva and liver punctuate by column switching for drug monitoring studies 182

Albumin
Procedure for the isolation of a large peptic fragment of human serum albumin 13

Alkaloids
Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 83

Amino acids
Glutamate and glutamine in cerebrospinal fluid and serum from healthy volunteers — analytical aspects 378

Amino peptidase-P
HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 129

3-(4′-Aminophenyl)-3-ethyl-1-methylpyrrolidine-2,5-dione
HPLC determination of imide inhibitors of aromatase in biological samples 163

3-(4′-Aminophenyl)-3-ethylpyrrolidine-2,5-dione
HPLC determination of imide inhibitors of aromatase in biological samples 163

21-Aminosteroids
Irregular retention properties of 21-aminosteroid antioxidants in octylsilane bonded-phase HPLC 285

21-Aminosteroids
HPLC determination of a 21-aminosteroid antioxidant in plasma 293

Anabolics
Identification and quantitation of trenbolone in bovine tissue by GC–MS 219

Anions
Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 29

Antibiotics
Determination of β-lactams and their biosynthetic intermediates in fermentation media by pre-column derivatisation followed by fluorescence detection 325

Antioxidants
Irregular retention properties of 21-aminosteroid antioxidants in octylsilane bonded-phase HPLC 285

Antioxidants
HPLC determination of a 21-aminosteroid antioxidant in plasma 293

Aromatase inhibitors
HPLC determination of imide inhibitors of aromatase in biological samples 163

Articaine
HPLC and preliminary pharmacokinetics of articaine and its 2-carboxy metabolite in human serum and urine 440

Cadralazine
HPLC determination of cadralazine in human plasma and urine 119

2-Carboxyarticaine
HPLC and preliminary pharmacokinetics of articaine and its 2-carboxy metabolite in human serum and urine 440

Carboxylic acids
New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 357

Catecholamines
Routine determination of urinary free catecholamines by HPLC with electrochemical detection 263

Catecholamines
Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 273

Catecholamines
LC determination of urinary catecholamines after one-step alumina extraction 373

Cefotaxime
HPLC assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 111

Ceftriaxone
HPLC assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 111
Cephalosporins
HPLC assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 111
Cephalosporins
Determination of β-lactams and their biosynthetic intermediates in fermentation media by pre-column derivatisation followed by fluorescence detection 325
Cisapride
Determination of cisapride in plasma and animal tissues by HPLC 195
Citrate
New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 357
Copper(II) diethyldithiocarbamate
Determination of disulfiram and two of its metabolites in urine by reversed-phase LC and spectrophotometric detection after post-column complexation 95
Cysteine
Measurement of plasma thiols after derivatization with monobromobimane 141
Deacetylthymoxamine
HPLC assay of the metabolites of thymoxamine 205
7-Deoxydoxorubicin aglycone
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
7-Deoxydoxorubicinol aglycone
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
Deoxyspergualin
Determination of deoxyspergualin by GC–selected-ion monitoring 39
N-Depropylpropafenone
HPLC separation and MS identification of propafenone, 5-hydroxypropafenone and N-depropylpropafenone 211
Desacetylcefotaxime
HPLC assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 111
Desacetylvinblastine
Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 83
Desmethyladinazolam
HPLC assay for adinazolam and its desmethyl metabolite 215
N-Desmethyldiazepam
Improved method for the determination of diazepam and N-desmethyldiazepam in plasma using capillary GC and nitrogen–phosphorus detection 398
Diazepam
Improved method for the determination of diazepam and N-desmethyldiazepam in plasma using capillary GC and nitrogen–phosphorus detection 398
Diethyldithiocarbamate
Determination of disulfiram and two of its metabolites in urine by reversed-phase LC and spectrophotometric detection after post-column complexation 95
3,4-Dihydroxyphenylacetic acid
Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 273
Dimercaptosuccinic acid
GC analysis of urinary dimercaptosuccinic acid 231
N^4-[2,6-Dimethoxy-4-methyl-5-[(3-trifluoromethyl)phenoxy]-8-quinolinyl]-1,4-pentanediamine
Rapid and sensitive quantitative analysis of the new antimalarial N^4-[2,6-dimethoxy-4-methyl-5-[(3-trifluoromethyl)phenoxy]-8-quinolinyl]-1,4-pentanediamine in plasma by LC and electrochemical detection 347
Dipeptidyl peptidase IV
HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 129
Dipipanone
Measurement of dipipanone using capillary GC 152
Disopyramide
Simultaneous assay of disopyramide and monodesisopropyldisopyramide enantiomers in biological samples by LC 424
Disulfiram
Determination of disulfiram and two of its metabolites in urine by reversed-phase LC and spectrophotometric detection after post-column complexation 95
Dopamine
Routine determination of urinary free catecholamines by HPLC with electrochemical detection 263

Dopamine
LC determination of urinary catecholamines after one-step alumina extraction 373
Doxorubicin
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
Doxorubicin aglycone
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
Doxorubicinol
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
Doxorubicinol aglycone
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
Enantiomer separation
Determination of drug enantiomers in biological samples by coupled column LC and LC–MS 61
Enzymes
HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 129
Enzymes
Determination of phenol sulphotransferase activity by HPLC 136
Enzymes
Affinity chromatography of human thrombin on modified silica 385
Enzymes
Highly sensitive assay for xanthine oxidase activity by HPLC with fluorescence detection 392
4′-Epidoxorubicin
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
4′-Epidoxorubicin glucuronide
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
4′-Epidoxorubicinol
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
4′-Epidoxorubicinol glucuronide
New method for the determination of doxorubicin, 4′-epidoxorubicin and all known metabolites in cardiac tissue 103
Epinephrine
Routine determination of urinary free catecholamines by HPLC with electrochemical detection 263
Epinephrine
LC determination of urinary catecholamines after one-step alumina extraction 373
Esmolol
Analysis of esmolol in human blood by HPLC and its application to pharmacokinetic studies 430
Fumarate
New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 357
Furprofen
Determination of furprofen in human plasma by solid-phase extraction and reversed-phase HPLC 201
Gasoline
GC determination for forensic purposes of petroleum fuel inhaled just before fatal burning 49
Glutamate
Glutamate and glutamine in cerebrospinal fluid and serum from healthy volunteers — analytical aspects 378
Glutamine
Glutamate and glutamine in cerebrospinal fluid and serum from healthy volunteers – analytical aspects 378
Glutathione
Measurement of plasma thiols after derivatization with monobromobimane 141
High-capacity immunoaffinity sorbents
Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their $F_{ab'}$ fragments for isolation of proteins 243
Homocysteine
Measurement of plasma thiols after derivatization with monobromobimane 141
Homovanillic acid
Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 273
Human serum albumin
Procedure for the isolation of a large peptic fragment of human serum albumin 13

Hydrocarbons
GC determination for forensic purposes of petroleum fuel inhaled just before fatal burning 49
4-Hydroxydebrisoquine
HPLC quantitation of 4-hydroxydebrisoquine in microsomal incubates by use of silica columns, aqueous mobile phase and automated column switching 435
5-Hydroxyindole-3-acetic acid
Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 273
4-Hydroxy-3-methoxymandelic acid
Automatic determination of urinary 4-hydroxy-3-methoxymandelic (vanillylmandelic) acid by LC with electrochemical detection 255
5-Hydroxypropafenone
HPLC separation and MS identification of propafenone, 5-hydroxypropafenone and N-depropylpropafenone 211
Imide inhibitors of aromatase
HPLC determination of imide inhibitors of aromatase in biological samples 163
Immunoaffinity sorbents
Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their $F_{ab'}$ fragments for isolation of proteins 243
Immunoglobulin $F_{ab'}$ fragments
Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their $F_{ab'}$ fragments for isolation of proteins 243
Inorganic anions
Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 29
Isotretinoin
Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 303
Janus Green
Use of the dye Janus Green to facilitate quantitative analysis of phosphoinositides 445
Kerosene
GC determination for forensic purposes of petroleum fuel inhaled just before fatal burning 49
α-Keto acids
Determination of α-keto acids in rat hindquarter muscle by HPLC with fluorescence detection 361
α-Ketocaproate
Determination of α-keto acids in rat hindquarter muscle by HPLC with fluorescence detection 361
α-Ketoisocaproate
Determination of α-keto acids in rat hindquarter muscle by HPLC with fluorescence detection 361
α-Ketoisovalerate
Determination of α-keto acids in rat hindquarter muscle by HPLC with fluorescence detection 361
α-Keto-β-methylvalerate
Determination of α-keto acids in rat hindquarter muscle by HPLC with fluorescence detection 361
15-Ketoprostaglandin $F_{2\alpha}$
Dimethylisopropylsilyl ether derivatives in GC–MS of some prostanoids. Application to the microdetermination of 15-keto-prostaglandin $F_{2\alpha}$ formed by enzymatic conversion of $PGF_{2\alpha}$ in vitro 1
Krebs cycle acids
New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 357
β-Lactams
Determination of β-lactams and their biosynthetic intermediates in fermentation media by pre-column derivatisation followed by fluorescence detection 325
Leucovorin
Direct resolution of the stereoisomers of leucovorin and 5-methyltetrahydrofolate using a bovine serum albumin HPLC chiral stationary phase coupled to an achiral phenyl column 158
Malate
New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 357
Methylergonovine
HPLC method for the determination of methysergide and methylergonovine in human plasma 416
N^1-Methyl-2-pyridone-5-carboxamide
Simultaneous micro-determination of nic-

otinamide and its major metabolites, N^1-methyl-2-pyridone-5-carboxamide and N^1-methyl-4-pyridone-3-carboxamide, by HPLC 23

N^1-Methyl-4-pyridone-3-carboxamide
Simultaneous micro-determination of nicotinamide and its major metabolites, N^1-methyl-2-pyridone-5-carboxamide and N^1-methyl-4-pyridone-3-carboxamide by HPLC 23

5-Methyltetrahydrofolate
Direct resolution of the stereoisomers of leucovorin and 5-methyltetrahydrofolate using a bovine serum albumin HPLC chiral stationary phase coupled to an achiral phenyl column 158

Methysergide
HPLC method for the determination of methysergide and methylergonovine in human plasma 416

Mitoxantrone
Direct determination of mitoxantrone and its mono- and dicarboxylic metabolites in plasma and urine by HPLC 337

Monobromobimane
Measurement of plasma thiols after derivatization with monobromobimane 141

Monodesisopropyldisopyramide
Simultaneous assay of disopyramide and monodesisopropyldisopyramide enantiomers in biological samples by LC 424

N-Monomethyldeacetylthymoxamine
HPLC assay of the metabolites of thymoxamine 205

3-O-Morpholinoethylmorphine
Determination of pholcodine in biological fluids by HPLC with fluorescence detection 170

Mucins
Sulphate quantification in rabbit tracheobronchial mucus by reversed-phase ion-pair chromatography 147

Nicotinamide
Simultaneous micro-determination of nicotinamide and its major metabolites, N^1-methyl-2-pyridone-5-carboxamide and N^1-methyl-4-pyridone-3-carboxamide, by HPLC 23

Nitrazepam
Sensitive GC method for determining nitrazepam in serum and saliva 403

Nitrite
Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 29

Norepinephrine
Routine determination of urinary free catecholamines by HPLC with electrochemical detection 263

Norepinephrine
LC determination of urinary catecholamines after one-step alumina extraction 373

4-Oxoisotretinoin
Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 303

4-Oxotretinoin
Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 303

Penicillins
Determination of β-lactams and their biosynthetic intermediates in fermentation media by pre-column derivatization followed by fluorescence detection 325

Peptides
Procedure for the isolation of a large peptic fragment of human serum albumin 13

Peptides
HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 129

Perchloric acid extracts
New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 357

Petroleum fuel
GC determination for forensic purposes of petroleum fuel inhaled just before fatal burning 49

Phenol sulphotransferase
Determination of phenol sulphotransferase activity by HPLC 136

Pholcodine
Determination of pholcodine in biological fluids by HPLC with fluorescence detection 170

Phosphatidic acid
Sensitive HPLC method for the determination of phosphatidic acid 367

Phosphoinositides
Use of the dye Janus Green to facilitate quantitative analysis of phosphoinositides 445
Piperacillin
HPLC assay for piperacillin in aqueous humor of the eye 177
Proline
HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 129
Propafenone
HPLC separation and MS identification of propafenone, 5-hydroxypropafenone and N-depropylpropafenone 211
Propranolol
Separation of enantiomers of derivatised or underivatised propranolol by means of HPLC 189
Prostanoids
Dimethylisopropylsilyl ether derivatives in GC–MS of some prostanoids. Application to the microdetermination of 15-ketoprostaglandin $F_{2\alpha}$ formed by enzymatic conversion of $PGF_{2\alpha}$ in vitro 1
Pseudoephedrine
Sensitive quantification of pseudoephedrine in human plasma and urine by HPLC 73
Pyridoxal
Simple isocratic HPLC method for the separation of the six vitamers of vitamin B_6 315
Pyridoxal 5′-phosphate
Simple isocratic HPLC method for the separation of the six vitamers of vitamin B_6 315
Pyridoxamine
Simple isocratic HPLC method for the separation of the six vitamers of vitamin B_6 315
Pyridoxamine 5′-phosphate
Simple isocratic HPLC method for the separation of the six vitamers of vitamin B_6 315
Pyridoxine
Simple isocratic HPLC method for the separation of the six vitamers of vitamin B_6 315
Pyridoxine 5′-phosphate
Simple isocratic HPLC method for the separation of the six vitamers of vitamin B_6 315
Retinoids
Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 303
Serotonin
Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 273
Steroids
Identification and quantitation of trenbolone in bovine tissue by GC–MS 219
Steroids
Irregular retention properties of 21-aminosteroid antioxidants in octylsilane bonded-phase HPLC 285
Steroids
HPLC determination of a 21-aminosteroid antioxidant in plasma 293
Succinate
New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 357
Sulphate
Sulphate quantification in rabbit tracheobronchial mucus by reversed-phase ion-pair chromatography 147
Sulphide
Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 29
Terbutaline
Determination of drug enantiomers in biological samples by coupled column LC and LC–MS 61
Tetraprenylacetone
Determination of tetraprenylacetone in human plasma by HPLC with fluorescence derivatization using dansylhydrazine 410
Thiocyanate
Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 29
Thiols
Measurement of plasma thiols after derivatization with monobromobimane 141

Thiosulphate
Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 29
Thrombin
Affinity chromatography of human thrombin on modified silica 385
Thymoxamine
HPLC assay of the metabolites of thymoxamine 205
Trenbolone
Identification and quantitation of trenbolone in bovine tissue by GC–MS 219
Tretinoin
Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 303
Vanillylmandelic acid
Automatic determination of urinary 4-hydroxy-3-methoxymandelic (vanillylmandelic) acid by LC with electrochemical detection 255
Vanillylmandelic acid
Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 273
Vinblastine
Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 83
Vinca alkaloids
Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 83
Vincristine
Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 83
Vindesine
Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 83
Vitamin B
Simultaneous micro-determination of nicotinamide and its major metabolites, N^1-methyl-2-pyridone-5-carboxamide and N^1-methyl-4-pyridone-3-carboxamide, by HPLC 23
Vitamin B_6
Simple isocratic HPLC method for the separation of the six vitamers of vitamin B_6 315
Xanthine oxidase
Highly sensitive assay for xanthine oxidase activity by HPLC with fluorescence detection 392